

Factorization method for second order functional equations

Tomasz Goliński¹, Anatol Odziejewicz²

Institute of Theoretical Physics
University in Białystok
Lipowa 41, 15-424 Białystok, Poland

PACS: 02.30.Ks, 02.30.Vv

CONTENTS

Introduction	1
1. Calculus generated by the bijection of subset of the real numbers	2
2. Hilbert spaces with scalar product given by τ -integral	4
3. Factorization method	6
4. Change of variables	9
5. Some examples	10
5.1. Particular solution of (3.9)	10
5.2. q -Hahn polynomials	12
5.3. The case $g_k = \text{const}$ and $\tau(x) = qx$	13
5.4. The case $g_k = \text{const}$ and τ is a fractional map	15
Appendix A. τ -Riccati equation and τ -Darboux transform	17
Acknowledgement	21
References	22

INTRODUCTION

In this paper we develop the theory of equations of the form

$$(0.1) \quad \alpha(x)\psi(\tau(x)) + \beta(x)\psi(x) + \gamma(x)\psi(\tau^{-1}(x)) = \lambda\psi(x)$$

generated by a bijection $\tau : X \rightarrow X$ of the real line subset $X \subset \mathbb{R}$.

The equations of this type can be regarded as an alternate discretization of the second order differential equations including Schrödinger equation and generalization of difference and q -difference equations. They also emerge from the change of variables in difference equations. On the other hand the functional equations by themselves are of interest and have many important applications. They have been investigated in many monographs and papers from the point of view of functional analysis, especially C^* -algebras methods, see e.g. [2, 3]. We hope that our approach can be applied also in the numerical analysis of some problems.

Our approach is based on τ -difference and τ -integral calculus, see e.g. [11], which is a direct generalization of the standard one and is ideologically related to the analysis on time scales. This gives us a possibility to generalize the factorization method elaborated effectively by many authors in the differential case, e.g. see [4, 10]. Our approach allows one to consider this method in q -case, see [5] for

¹tomaszg@alpha.uwb.edu.pl

²aodziejew@labfiz.uwb.edu.pl

application, and due to generality of τ has the benefit that it is equivariant under the change of variables.

Section 1 defines basic notions and operators of τ -calculus as well as formulates some relations between them. In Section 2 we introduce and study Hilbert spaces with the scalar product given by τ -integral on the orbits of the semigroup of natural numbers $(\mathbb{N} \cup \{0\}, +)$ or group of integer numbers $(\mathbb{Z}, +)$ acting on X by the iterations of the bijection τ . Then, in Section 3 we construct the chain of eigenproblems in these Hilbert spaces by the application of the factorization method. This construction gives a possibility to solve the eigenproblems for the chain of Hamiltonians if one knows a solution to the initial one.

In Section 4 we discuss the change of variables and induced transformation of Hilbert spaces, eigenproblems and parametrizing functions. Some remarks about equivalence of one τ -difference operator to another are also presented.

In Section 5 we show that our method includes the one investigated in the theory of q -Hahn orthogonal polynomials, which are q -deformation of the classical ones, see [12]. In this section we also present some examples.

Some general facts about τ -equations is presented in Appendix A, i.e. some formulas for solutions. Reader can find there easy results which are the generalization of theorems about q -Riccati equation from [13].

Let us stress finally the importance of the functional equation (0.1) in the theory of discrete and q -discrete polynomials, see [12]. This link is also important to the integration of some quantum optical systems, see [9].

1. CALCULUS GENERATED BY THE BIJECTION OF SUBSET OF THE REAL NUMBERS

Let us introduce preliminary definitions and facts of the calculus generated by the bijection $\tau : X \rightarrow X$ of the subset $X \subset \mathbb{R}$ (see also [11]). Let $\mathcal{F}(X)$ denotes the algebra of all complex valued functions $\varphi : X \rightarrow \mathbb{C}$. Let $\mathcal{O}(x) = \{\tau^n(x) | n \in \mathbb{Z}\}$ be the orbit of point x under action of group $\mathbb{Z}$ given by τ and $\mathcal{O}^+(x) = \{\tau^n(x) | n \in \mathbb{N} \cup \{0\}\}$ — orbit under action of semigroup $\mathbb{N} \cup \{0\}$.

The subject of the paper is linked to the investigation of the following linear operators

$$(1.1) \quad T\varphi(x) := \varphi(\tau(x))$$

$$(1.2) \quad M_f\varphi(x) := f(x)\varphi(x)$$

$$(1.3) \quad \partial_\tau\varphi(x) := \frac{\varphi(x) - \varphi(\tau(x))}{x - \tau(x)},$$

where $f \in \mathcal{F}(X)$.

The operator ∂_τ is a direct generalization of q -derivative ∂_q , which corresponds to the special case when $\tau(x) = qx$. We shall call it **τ -derivative**. It satisfies the following analogue of Leibniz rule

$$(1.4) \quad \partial_\tau(\varphi\psi) = (T\varphi)\partial_\tau\psi + \psi\partial_\tau\varphi.$$

We define τ -integral as the solution of the equation

$$(1.5) \quad \partial_\tau\varphi = \psi$$

for given $\psi \in \mathcal{F}(X)$. Applying the operator T^n to both sides of (1.5) one obtains

$$(1.6) \quad \varphi(\tau^n(x)) - \varphi(\tau^{n+1}(x)) = (\tau^n(x) - \tau^{n+1}(x))\psi(\tau^n(x))$$

for $n = 0, 1, 2, \dots$, where $\tau^n = \underbrace{\tau \circ \dots \circ \tau}_{n \text{ times}}$ and thus

$$(1.7) \quad \varphi(x) - \varphi(\tau^\infty(x)) = \sum_{n=0}^{\infty} (\tau^n(x) - \tau^{n+1}(x))\psi(\tau^n(x)).$$

The formula (1.7) has sense only if $\tau^n(x)$ has the limit when $n \rightarrow \infty$ and φ is continuous at $\tau^\infty(x) \in X$. Hence, if these conditions are satisfied, one can define the **τ -integral**:

$$(1.8) \quad \int_{\tau^\infty(x)}^x \psi(t) d_\tau t = \int_{\mathcal{O}^+(x)} \psi(t) d_\tau t := \sum_{n=0}^{\infty} (\tau^n(x) - \tau^{n+1}(x))\psi(\tau^n(x)).$$

Let $[a, b] \subset X$ be an interval such that $\tau^\infty(a) = \tau^\infty(b)$ then we shall define the integral of ψ over $[a, b]$ by:

$$(1.9) \quad \int_a^b \psi(t) d_\tau t = \int_{\mathcal{O}^+(a) \cup \mathcal{O}^+(b)} \psi(t) d_\tau t := \int_{\tau^\infty(b)}^b \psi(t) d_\tau t - \int_{\tau^\infty(a)}^a \psi(t) d_\tau t.$$

Let us note that integral is only formally over $[a, b]$ and it is based on $\mathcal{O}^+(a) \cup \mathcal{O}^+(b)$. Similarly we will define τ -integral over $\mathcal{O}(x)$ by

$$(1.10) \quad \int_{\mathcal{O}(x)} \psi(t) d_\tau t := \sum_{n=-\infty}^{\infty} (\tau^n(x) - \tau^{n+1}(x))\psi(\tau^n(x)).$$

The terminology introduced above is justified by the following properties of the τ -integral:

$$(1.11) \quad \int_a^b (\partial_\tau \psi)(t) d_\tau t = \psi(b) - \psi(a),$$

$$(1.12) \quad \partial_\tau \int_{\tau^\infty(x)}^x \psi(t) d_\tau t = \psi(x).$$

Moreover we have the following formula for the change of variables

$$(1.13) \quad \int_a^b (T\psi)(t) \rho(t) d_\tau t = \int_{\tau(a)}^{\tau(b)} \psi(t) (\partial_\tau \tau^{-1})(t) (T^{-1}\rho)(t) d_\tau t.$$

Let us denote by $\mathcal{D}(X) \subset \mathcal{F}(X)$ the subalgebra of functions with finite support. This subalgebra is preserved by T and ∂_τ , as well as by M_f .

For affine bijection $\tau(x) = qx + h$ all formulae above reduce to the corresponding formulae of q -difference ($h = 0$) or h -difference ($q = 1$) calculus. If $q \rightarrow 1$ ($h = 0$) or $h \rightarrow 0$ ($q = 1$) one obtains the formulae of the standard differential and integral calculus. Formulae (1.11) and (1.12) become then the fundamental theorem of calculus and (1.13) reduces to identity.

2. HILBERT SPACES WITH SCALAR PRODUCT GIVEN BY τ -INTEGRAL

Let us fix some weight function $\rho : X \rightarrow \mathbb{R}$. We can define the scalar product by the τ -integral over the set $\mathcal{S}$ by the formula

$$(2.1) \quad \langle \psi | \varphi \rangle := \int_{\mathcal{S}} \overline{\psi}(x) \varphi(x) \rho(x) d_{\tau} x,$$

where $\mathcal{S}$ is one of the following sets $\mathcal{O}(x_0)$, $\mathcal{O}^+(a)$, $\mathcal{O}^+(a) \cup \mathcal{O}^+(b)$ and φ, ψ are complex valued functions defined on X .

For the positivity of (2.1) we shall assume that $(\tau^n(b) - \tau^{n+1}(b))\rho(\tau^n(b)) \geq 0$ and $(\tau^n(a) - \tau^{n+1}(a))\rho(\tau^n(a)) \leq 0$, $n \in \mathbb{N} \cup \{0\}$ for semigroup orbits case or $(\tau^n(x_0) - \tau^{n+1}(x_0))\rho(\tau^n(x_0)) \geq 0$ for group orbit case.

Let us denote by $L^2(X, \rho d_{\tau})$ the Hilbert space of square-integrable functions in the sense of (2.1). It is clear that $\mathcal{D}(X) \subset L^2(X, \rho d_{\tau})$ for any weight function ρ . One shall keep in mind that elements of this space are classes of equivalence of functions from $\mathcal{F}(X)$, which can be identified with functions on $\mathcal{S}$.

If we consider semigroup orbit case $\mathcal{O}^+(a) \cup \mathcal{O}^+(b)$ and let $b = \tau^N(a)$ for some $N \in \mathbb{N}$, then $L^2(X, \rho d_{\tau})$ is finite dimensional. This case will not be discussed in the paper. So, it will be assumed everywhere that $b \neq \tau^N(a)$ for all $N \in \mathbb{N}$.

If one admits the case $\rho(x) = 0$ in (2.1) for some $x \in \mathcal{S}$, then the Hilbert space reduces to direct sum of Hilbert spaces realized by the functions on orbits or finite sets. So, one comes back to the cases defined above. The case of two semigroup orbits is also direct sum and it is considered here only due to the analogy with q -Hahn polynomials.

Let us now consider the operator T acting in Hilbert space $L^2(X, \rho d_{\tau})$ defined on $\mathcal{D}(X)$ by (1.1). Note that $\mathcal{D}(X)$ is dense in $L^2(X, \rho d_{\tau})$.

Proposition 2.1. *The adjoint operator T^* is the weighted shift operator given by:*

$$(2.2a) \quad (T^* \varphi)(x) = \begin{cases} \mu(x) \varphi(\tau^{-1}(x)) & \text{if } x \neq a \text{ and } x \neq b \\ 0 & \text{if } x = a \text{ or } x = b \end{cases}$$

for the semigroup orbits case and

$$(2.2b) \quad (T^* \varphi)(x) = \mu(x) \varphi(\tau^{-1}(x)),$$

for the group orbit case, where $\varphi \in \mathcal{D}_{T^*}$ and

$$(2.3) \quad \mu(x) := \partial_{\tau}(\tau^{-1})(x) \frac{\rho(\tau^{-1}(x))}{\rho(x)}.$$

The domain $\mathcal{D}_{T^*}$ of T^* contains $\mathcal{D}(X)$.

Let us mention the following properties of the operators T and T^* . For the group orbit case one has identity

$$(2.4a) \quad T^* T \varphi = \mu \varphi.$$

For the semigroup orbits case one obtains

$$(2.4b) \quad T^* T \varphi = \mu(1 - \chi_{\{a, b\}}) \varphi,$$

where $\chi_{\{a, b\}}$ is the characteristic function of the subset $\{a, b\} \subset X$ and thus $M_{\chi_{\{a, b\}}}$ is a projector onto this subset. For both group and semigroup orbits cases one has

$$(2.4c) \quad T T^* \varphi = (\mu \circ \tau) \varphi.$$

The operator T is bounded if and only if $\sup_{x \in \mathcal{S}} |\mu(x)| < \infty$ and then $\|T\| = \|T^*\| = (\sup_{x \in \mathcal{S}} |\mu(x)|)^{\frac{1}{2}}$.

Let χ_a and χ_b be the characteristic functions (indicators) of $\mathcal{O}^+(a)$ and $\mathcal{O}^+(b)$. The orthogonal projectors M_{χ_a} and M_{χ_b} map $\mathcal{D}(X)$ into itself. Since operator T preserves subspaces $L^2(\mathcal{O}^+(a), \rho d_\tau)$ and $L^2(\mathcal{O}^+(b), \rho d_\tau)$, it can be reduced to the components of the partition

$$(2.5) \quad L^2(X, \rho d_\tau) = L^2(\mathcal{O}^+(a), \rho d_\tau) \oplus L^2(\mathcal{O}^+(b), \rho d_\tau)$$

in the sense of [1]. Thus we can consider each orbit separately.

Let us now consider the sequence of Hilbert spaces $\mathcal{H}_k := L^2(X, \rho_k d_\tau)$ with the weight functions ρ_k , $k \in \mathbb{N} \cup \{0\}$ related by the recurrences

$$(2.6) \quad \rho_{k+1} = \eta_k \rho_k$$

and

$$(2.7) \quad \rho_{k+1} = T(B_k \rho_k)$$

for $B_k, \eta_k \in \mathcal{F}(X)$. Since we demand two distinct formulae for ρ_{k+1} , we need to add the consistency condition on η_k and B_k , namely

$$(2.8) \quad T(B_k \rho_k) = \eta_k \rho_k$$

and in semigroup case we additionally impose a boundary conditions

$$(2.9) \quad B_k(a) \rho_k(a) = B_k(b) \rho_k(b) = 0$$

for any $k \in \mathbb{N} \cup \{0\}$.

Let us denote by g_k the ratio of B_k and B_{k+1} :

$$(2.10) \quad B_{k+1} = g_k B_k.$$

Then we can easily see that formulas (2.6)–(2.9) are automatically satisfied if (2.8)–(2.9) are valid for $k = 0$ and

$$(2.11) \quad \eta_{k+1} = T(g_k \eta_k)$$

for any $k \in \mathbb{N} \cup \{0\}$.

By introducing the function

$$(2.12) \quad A_k(x) = \frac{B_k(x) - \eta_k(x)}{x - \tau(x)}$$

and expressing η_k by A_k and B_k , one can rewrite equation (2.8) in the form

$$(2.13) \quad \partial_\tau(B_k \rho_k) = A_k \rho_k$$

In the case of $\tau(x) = qx$ and when $q \rightarrow 1$, the equation (2.13) corresponds to the well known Pearson equation of the theory of classical orthogonal polynomials. This motivates us to call the consistency equation (2.8) the τ -**Pearson** equation.

We shall now consider the operators defined formally by (1.1)–(1.3) as operators in Hilbert spaces. For $f \in \mathcal{F}(X)$ we shall consider the operators of multiplication by the function f as

$$(2.14) \quad M_f : \mathcal{H}_k \rightarrow \mathcal{H}_{k+1}.$$

The operator T acts in each of the spaces separately

$$(2.15) \quad T : \mathcal{H}_k \rightarrow \mathcal{H}_k$$

and the τ -derivative becomes now the operator

$$(2.16) \quad \partial_\tau = M_{(id-\tau)^{-1}}(1-T) : \mathcal{H}_k \rightarrow \mathcal{H}_{k+1}.$$

Since operators M_f , T and ∂_τ are in general unbounded, we take $\mathcal{D}(X)$ as their domain and then close them. We will denote their closure by the same symbols. We also omit indices k in symbols of operators M , T and ∂_τ to simplify the notation.

In these settings we can apply equation (2.8) to calculate explicitly the term μ_k appearing in formula (2.3) for T^* :

$$(2.17) \quad \mu_k(x) = \partial_\tau(\tau^{-1})(x) \frac{B_k(x)}{\eta_k(\tau^{-1}(x))}.$$

Proposition 2.2.

The adjoint operators $M_f^* : \mathcal{H}_{k+1} \rightarrow \mathcal{H}_k$ and $\partial_\tau^* : \mathcal{H}_{k+1} \rightarrow \mathcal{H}_k$, are given by

$$(2.18) \quad M_f^* = M_{\eta_k \bar{f}}$$

$$(2.19) \quad \partial_\tau^* = (1 - T^*)M_{\eta_k(id-\tau)^{-1}}.$$

The domains of M_f^* and ∂_τ^* contain $\mathcal{D}(X)$.

Proof: By means of (2.6) and (2.1) we get

$$\begin{aligned} \langle f\varphi|\psi\rangle_{k+1} &= \int_S \overline{f(x)\varphi(x)}\psi(x)\rho_{k+1}(x)d_\tau x = \\ &= \int_S \overline{\varphi(x)}\psi(x)\overline{f(x)}\eta_k(x)\rho_k(x)d_\tau x = \langle \varphi|\bar{f}\eta_k\psi\rangle_k \end{aligned}$$

for $\varphi, \psi \in \mathcal{D}(S)$. Formula for ∂_τ^* follows directly from (2.18) and definition of ∂_τ (2.16). $\square$

3. FACTORIZATION METHOD

The proposed method is the direct generalization of method for the second order differential equations, see [4, 10]. Instead of the equation of the form

$$(3.1) \quad \alpha(x)\psi(\tau(x)) + \beta(x)\psi(x) + \gamma(x)\psi(\tau^{-1}(x)) = \lambda\psi(x),$$

where $\lambda \in \mathbb{C}$ and $\alpha, \beta, \gamma \in \mathcal{F}(X)$, we will consider the sequence (chain) of the factorized equations

$$(3.2) \quad (\mathbf{A}_k^* \mathbf{A}_k \psi_k)(x) = \lambda_k \psi_k(x)$$

on $\psi_k \in \mathcal{H}_k$ with the operators $\mathbf{A}_k : \mathcal{H}_k \rightarrow \mathcal{H}_{k+1}$ and their adjoints $\mathbf{A}_k^* : \mathcal{H}_{k+1} \rightarrow \mathcal{H}_k$ given by

$$(3.3a) \quad \mathbf{A}_k := M_{h_k} \partial_\tau + M_{f_k}$$

$$(3.3b) \quad \mathbf{A}_k^* = (1 - T^*)M_{h_k \eta_k (id-\tau)^{-1}} + M_{\eta_k f_k},$$

where $f_k : X \rightarrow \mathbb{R}$, $h_k : X \rightarrow \mathbb{R}$ and $k \in \mathbb{N} \cup \{0\}$. We assume that $\mathbf{A}_k$ is a closure of the operator defined by (3.3a) on $\mathcal{D}(X)$ and $\mathbf{A}_k^*$ is automatically closed. Then by von Neumann theorem (see [1]) $\mathbf{A}_k^* \mathbf{A}_k$ is self-adjoint and thus closed. The essence of the approach is to postulate the relation

$$(3.4) \quad \mathbf{A}_k \mathbf{A}_k^* = d_k \mathbf{A}_{k+1}^* \mathbf{A}_{k+1} + c_k,$$

where $c_k, d_k \in \mathbb{C}$ and $k \in \mathbb{N} \cup \{0\}$. It follows from (3.4) that given a solution $\psi_k \in \mathcal{H}_k$ of equation (3.2), the function

$$(3.5) \quad \psi_{k+1} := \mathbf{A}_k \psi_k \in \mathcal{H}_{k+1}$$

is a solution of (3.2) for $k+1$ with $\lambda_{k+1} = \frac{\lambda_k - c_k}{d_k}$. So, starting from the equation (3.2) for $k=0$ and its solution ψ_0 , one generates the family of equations (3.2) with solutions given by

$$(3.6) \quad \psi_k = \prod_{i=0}^{k-1} \mathbf{A}_i \psi_0.$$

The operators $\mathbf{A}_k$ and $\mathbf{A}_k^*$ depend on the functions f_k, h_k as well as on η_k and B_k related by the recurrences (2.11) and (2.10) respectively.

Let us introduce the function

$$(3.7) \quad \varphi_k(x) := f_k(x) + \frac{h_k(x)}{x - \tau(x)},$$

which we will use instead of f_k .

Proposition 3.1. *Condition (3.4) is equivalent to the transformation rule*

$$(3.8) \quad \varphi_{k+1} h_{k+1} = \frac{h_k}{d_k} T \left(\frac{\varphi_k}{g_k} \right)$$

and the system of equations

$$(2.9) \quad \frac{d_k g_k(x) B_k(x) (h_{k+1}(\tau^{-1}(x)))^2}{(x - \tau(x))(\tau^{-1}(x) - x)} - (\varphi_k(x))^2 \eta_k(x) + c_k = \frac{1}{d_k g_k(\tau(x))} \left(\frac{d_k g_k(\tau(x)) B_k(\tau(x)) (h_k(x))^2}{(\tau(x) - \tau^2(x))(x - \tau(x))} - (\varphi_k(\tau(x)))^2 \eta_k(\tau(x)) \frac{(h_k(x))^2}{(h_{k+1}(x))^2} \right)$$

on the functions $B_k, \eta_k, \varphi_k, g_k, h_k, h_{k+1}$ and constants c_k, d_k , where $k \in \mathbb{N} \cup \{0\}$.

Proof: We compute explicitly left and right hand sides of (3.4). By substituting the (3.3), (3.8) to (3.4) and applying (2.4b) one has

$$(3.10) \quad \begin{aligned} \mathbf{A}_k \mathbf{A}_k^* &= \frac{-h_k(x) \varphi_k(\tau(x)) \eta_k(\tau(x))}{x - \tau(x)} T + \frac{(h_k(x))^2 B_k(\tau(x))}{(\tau(x) - \tau^2(x))(x - \tau(x))} + \\ &+ (\varphi_k(x))^2 \eta_k(x) - \frac{\varphi_k(x) B_k(x) h_k(\tau^{-1}(x))}{x - \tau(x)} T^{-1} \end{aligned}$$

and

$$(3.11) \quad \begin{aligned} \mathbf{A}_{k+1}^* \mathbf{A}_{k+1} &= \frac{-\varphi_{k+1}(x) \eta_{k+1}(x) h_{k+1}(x)}{x - \tau(x)} T + \frac{B_{k+1}(x) (h_{k+1}(\tau^{-1}(x)))^2}{(x - \tau(x))(\tau^{-1}(x) - x)} + \\ &+ (\varphi_{k+1}(x))^2 \eta_{k+1}(x) - \frac{B_{k+1}(x) \varphi_{k+1}(\tau^{-1}(x)) h_{k+1}(\tau^{-1}(x))}{x - \tau(x)} T^{-1}. \end{aligned}$$

In (3.11) we have removed the projector $1 - \chi_{\{a,b\}}$ appearing in (2.4b) for the case of orbits of semigroups. It was multiplied by a function B_{k+1} , which by (2.9) is equal to zero at the points a and b anyway.

Substituting (3.10) and (3.11) into (3.4) and comparing coefficients in front of the operators T, id and T^{-1} , one obtains

$$(3.12) \quad h_k(x) \varphi_k(\tau(x)) \eta_k(\tau(x)) = d_k h_{k+1}(x) \varphi_{k+1}(x) \eta_{k+1}(x),$$

$$\begin{aligned}
& \frac{B_k(\tau(x))(h_k(x))^2}{(\tau(x) - \tau^2(x))(x - \tau(x))} + (\varphi_k(x))^2 \eta_k(x) = \\
(3.13) \quad & = d_k \left(\frac{B_{k+1}(x)(h_{k+1}(\tau^{-1}(x)))^2}{(x - \tau(x))(\tau^{-1}(x) - x)} + (\varphi_{k+1}(x))^2 \eta_{k+1}(x) \right) + c_k
\end{aligned}$$

and

$$(3.14) \quad \varphi_k(x)B(x)h_k(\tau^{-1}(x)) = d_k B_{k+1}(x)\varphi_{k+1}(\tau^{-1}(x))h_{k+1}(\tau^{-1}(x)).$$

Using the transformation rules (2.11) and (2.10) one sees that conditions (3.12) and (3.14) are equivalent. They give the transformation rule (3.8)

$$(3.15) \quad \varphi_{k+1}h_{k+1} = h_k T \left(\frac{1}{d_k g_k} \varphi_k \right)$$

and condition (3.13) can be written in the form (3.9). Domains of $\mathbf{A}_k \mathbf{A}_k^*$ and $\mathbf{A}_{k+1}^* \mathbf{A}_{k+1}$ coincide due to closeness. $\square$

System of equations (3.9) can be regarded as an equation for gauge functions g_k (defined by (2.11)-(2.10)) or transformation rule for h_k given in non explicit way. Check Example 5.1 for explicit formulae for g_k in the case $h_k \equiv 1$ and $c_k = 0$ as well as convergence conditions. In the case $\tau(x) = qx$ and $g_k = \text{const}$ solutions to (3.9) have been classified in [5].

If we have the solution of (3.9) (both h_{k+1} and g_k) one can express the functions $(B_{k+1}, \eta_{k+1}, f_{k+1})$ of the $(k+1)^{st}$ step in the terms of the previous functions (B_k, η_k, f_k) .

Functions B_0, η_0, φ_0 and λ_0 are related to initial equation (3.1) by

$$(3.16) \quad \alpha(x) = -\frac{\varphi_0(x)\eta_0(x)h_0(x)}{x - \tau(x)},$$

$$(3.17) \quad \beta(x) = (\varphi_0(x))^2 \eta_0(x) + \frac{B_0(x)h_0(\tau^{-1}(x))}{(x - \tau(x))(\tau^{-1}(x) - x)},$$

$$(3.18) \quad \gamma(x) = -\frac{B_0(x)\varphi_0(\tau^{-1}(x))h_0(\tau^{-1}(x))}{x - \tau(x)},$$

and

$$(3.19) \quad \lambda = \lambda_0.$$

The formulae inverse to (3.16)-(3.19) have the form

$$(3.20) \quad B_0(x) = \frac{(x - \tau(x))(\tau^{-1}(x) - x)}{(h_0(\tau^{-1}(x)))^2} \left(\beta(x) + (x - \tau(x))\alpha(x) \frac{\varphi_0(x)}{h_0(x)} \right),$$

$$(3.21) \quad \eta_0(x) = -\frac{(x - \tau(x))\alpha(x)}{\varphi_0(x)h_0(x)},$$

where φ_0 and h_0 have to be chosen in such way that their ratio satisfies the equation of the τ -Riccati form (A.11):

$$(3.22) \quad \frac{\varphi_0(\tau(x))}{h_0(\tau(x))} = \frac{-\frac{\gamma(\tau(x))}{x - \tau(x)} - \frac{\varphi_0(x)}{h_0(x)} \beta(\tau(x))}{\frac{\varphi_0(x)}{h_0(x)} \alpha(\tau(x))(\tau(x) - \tau^2(x))}.$$

By this approach we have reduced the problem of construction of the chain of Hamiltonians to finding the solution of (3.9) and choice of initial conditions is ruled by (3.22).

4. CHANGE OF VARIABLES

We are going to consider an invertible change of variables

$$(4.1) \quad \kappa : X \longrightarrow Y.$$

It is convenient to assume that κ is homeomorphism in order to preserve the limits. Such transform allows us to modify the function τ in considered equations. Let us note that in differential equation change of variables keeps the differentiation operators and only changes the direction of derivation and multiplies them by some function factor. In discrete case one obtains in fact different derivations.

Change of variables defines the map from $\mathcal{F}(Y)$ to $\mathcal{F}(X)$ by

$$(4.2) \quad K : \mathcal{F}(Y) \ni f \longrightarrow Kf = f \circ \kappa \in \mathcal{F}(X).$$

We can associate to T the operator

$$(4.3) \quad \tilde{T} = K^{-1}TK$$

acting as the shift by

$$(4.4) \quad \tilde{\tau} = \kappa \circ \tau \circ \kappa^{-1}$$

on $\mathcal{F}(Y)$. One calls functions τ and $\tilde{\tau}$ equivalent if they are related by (4.4) for some κ . Operator M_f , $f \in \mathcal{F}(X)$, transforms to

$$(4.5) \quad K^{-1}M_fK = M_{f \circ \kappa^{-1}}.$$

Thus similarity transformation $K^{-1} \cdot K$ maps shift operators to shift operators and multiplication operators to multiplication operators. However it turns out that derivation is preserved by this transform only up to the function factor:

$$(4.6) \quad K^{-1}\partial_\tau K = M_{(\partial_\tau \kappa) \circ \kappa^{-1}} \partial_{\tilde{\tau}}.$$

For example let $X = [0, \infty)$, $\tau(x) = qx$, $Y = \mathbb{R}$ for $0 < q \neq 1$. Then $\kappa(x) = \ln(x)$ transforms τ to

$$(4.7) \quad \tilde{\tau}(y) = y + \ln q$$

and

$$(4.8) \quad K\partial_q K^{-1} = M_{\frac{-\ln q}{(1-q)e^y}} \partial_{\tilde{\tau}}.$$

Thus we can map q -difference equation to h -difference equation although there may appear function factor changing the form of equation.

For given (X, τ) and $(Y, \tilde{\tau})$ in general it is not possible to construct such κ that (4.4) would be valid. It is the case for example for $X = Y = [0, 1]$, $\tau(x) = x^2$ and $\tilde{\tau}(y) = \frac{2}{5}y^3 - \frac{3}{5}y^2 + \frac{6}{5}y$. It follows from the fact that κ maps fixed points of τ into fixed points of $\tilde{\tau}$. Since τ has two fixed points (0 and 1) while $\tilde{\tau}$ has three (0, $\frac{1}{2}$ and 1), thus $\tilde{\tau}$ is not equivalent to τ . In some cases however one can restrict X and Y in such way that the function would become equivalent but it may happen that κ is not given in terms of elementary functions thus rendering further calculations extremely complicated.

Let us now consider change of variables defined by (4.1) from the point of view of Hilbert spaces and introduced operators and functions. Let us consider Hilbert

spaces $\tilde{\mathcal{H}}_k$ of functions on the set Y defined in such way that it makes K an unitary operator mapping $\mathcal{H}_k$ to $\tilde{\mathcal{H}}_k$. Simple calculations shows that we have to consider the spaces $\tilde{\mathcal{H}} = L^2(\kappa(X), \tilde{\rho}_k d\tilde{\tau})$ with weight given by

$$(4.9) \quad \tilde{\rho}_k = (\partial_{\tilde{\tau}} \kappa^{-1}) K^{-1} \rho_k.$$

In order to keep formulas for transformation of $\tilde{\rho}_k$ the same as (2.6) and (2.7) one has to define

$$(4.10) \quad \tilde{B}_k = \frac{\tilde{T}^{-1}(\partial_{\tilde{\tau}} \kappa^{-1})}{(\partial_{\tilde{\tau}} \kappa^{-1})} K^{-1} B_k$$

and

$$(4.11) \quad \tilde{\eta}_k = K^{-1} \eta_k.$$

These functions automatically satisfy $\tilde{\tau}$ -Pearson equation and obey the same transformation laws with function

$$(4.12) \quad \tilde{g}_k = K^{-1} g_k.$$

If we put the following transformation rules for functions f_k and h_k

$$(4.13) \quad \tilde{h}_k = K^{-1} (h_k \partial_{\tau}(\kappa))$$

$$(4.14) \quad \tilde{f}_k = K^{-1} f_k,$$

then operators $\tilde{\mathbf{A}}_k = K^{-1} \mathbf{A}_k K$ and $\tilde{\mathbf{A}}_k^* = K^{-1} \mathbf{A}_k^* K$ are of the same form as original with τ replaced by $\tilde{\tau}$. Clearly $\tilde{\mathbf{A}}_k^*$ is an adjoint of $\tilde{\mathbf{A}}_k$ since K is an unitary. Let us note that due to (4.13) the condition $h_k \equiv 1$ is not preserved by the change of variables except in the case of translation $\kappa(x) = x + c$, $c \in \mathbb{R}$.

Solutions of considered second order equation (3.2) transform as follows

$$(4.15) \quad \tilde{\psi}_k = K^{-1} \psi_k.$$

Thus we were able to map the chain of Hamiltonians on X to the chain of Hamiltonians on Y preserving all relations introduced in Section 2 and 3. This shows that considered method is equivariant with respect to the change of variables. It was not true for the case of q -difference equations.

5. SOME EXAMPLES

We are going to present several examples for the application of factorization method described above.

5.1. Particular solution of (3.9). We are going to construct a solution equation (3.9) in terms of τ -integrals by application of propositions from Appendix A.

We consider the simplest case $h_k \equiv 1$. Let us introduce the function

$$(5.1) \quad \xi_k(x) := (\varphi_k(x))^2 \eta_k(x) - \frac{d_k g_k(x) B_k(x)}{(x - \tau(x))(\tau^{-1}(x) - x)}.$$

After substituting ξ_k instead of g_k the equation (3.9) takes the form

$$(5.2) \quad \xi_k(x) = \frac{\left(\frac{B_k(\tau(x))}{(\tau(x) - \tau^2(x))(x - \tau(x))} - c_k \right) \xi_k(\tau(x)) + c_k (\varphi_k(\tau(x)))^2 \eta_k(\tau(x))}{(\varphi_k(\tau(x)))^2 \eta_k(\tau(x)) - \xi_k(\tau(x))},$$

which is also a τ -Riccati equation (A.11). It is more convenient than the form (3.9) because in the case of $c_k = 0$ one can apply the Proposition A.1 from Appendix

A which allows one to write down formulae for solutions. To that end we have to invert the fractional map on right-hand side of (5.2) and use it to define matrix $\mathbf{\Lambda}_k(x)$ appearing in (A.9). We get:

$$(5.3) \quad \mathbf{\Lambda}_k(x) = \begin{pmatrix} 1 & \frac{(x-\tau(x))(\tau(x)-\tau^2(x))}{B_k(\tau(x))} \\ 0 & \frac{(x-\tau(x))(\tau(x)-\tau^2(x))\varphi_k(\tau(x))^2\eta_k(\tau(x))}{B_k(\tau(x))} \end{pmatrix}.$$

Since this matrix can be singular at $\tau^\infty(x)$, we perform τ -Darboux transform (A.34) to $\mathbf{\Lambda}_k(x)$ with $\delta_2 = \lambda$ and $\delta_1 = 0$. In order to apply Proposition A.3 we have to compute $\tilde{\mathbf{\Lambda}}'_k$ defined by (A.8):

$$(5.4) \quad \tilde{\mathbf{\Lambda}}'_k(x) = \begin{pmatrix} 0 & -\frac{(\tau(x)-\tau^2(x))(x-\tau^\infty(x))^\lambda}{B_k(\tau(x))} \\ 0 & \frac{1}{x-\tau(x)} - \frac{\varphi_k(\tau(x))^2\eta_k(\tau(x))}{B_k(\tau(x))} \frac{(\tau(x)-\tau^2(x))(x-\tau^\infty(x))^\lambda}{(\tau(x)-\tau^\infty(x))^\lambda} \end{pmatrix}.$$

Let us put $\lambda = -1$. If the functions f_k , η_k and B_k are regular and non-vanishing at $\tau^\infty(x)$ and τ is differentiable at $\tau^\infty(x)$ then we have

$$(5.5) \quad \lim_{n \rightarrow \infty} \tilde{\mathbf{\Lambda}}'_k(\tau^n(x)) = \begin{pmatrix} 0 & -\left(\frac{d\tau}{dx}(\tau^\infty(x)) - \frac{d\tau^2}{dx}(\tau^\infty(x))\right) (B_k(\tau^\infty(x)))^{-1} \\ 0 & 0 \end{pmatrix}.$$

In such a way the singularity in $\tilde{\mathbf{\Lambda}}_k$ is removed. If we assume that τ satisfies the condition *i*) of Proposition A.3 then the resolvent matrix is well defined and by Proposition A.1 we find

$$(5.6) \quad \mathbf{\Lambda}'_{k\infty}(x) = \begin{pmatrix} 1 & F_k(x) \\ 0 & \exp\left(\int_{\tau^\infty(x)}^x \frac{\ln \frac{(\tau(t)-\tau^\infty(t))(t-\tau(t))(\tau(t)-\tau^2(t))\varphi_k(\tau(t))^2\eta_k(\tau(t))}{(t-\tau^\infty(t))B_k(\tau(t))}}{t-\tau(t)} d_\tau t\right) \end{pmatrix},$$

where $F_k(x)$ is given by

$$\begin{aligned} F_k(x) = & \exp\left(\int_{\tau^\infty(x)}^x \frac{\ln \frac{(\tau(t)-\tau^\infty(t))(t-\tau(t))(\tau(t)-\tau^2(t))\varphi_k(\tau(t))^2\eta_k(\tau(t))}{(t-\tau^\infty(t))B_k(\tau(t))}}{t-\tau(t)} d_\tau t\right) \times \\ & \times \int_{\tau^\infty(x)}^x \frac{(\tau(t)-\tau^2(t))}{(t-\tau^\infty(t))B_k(\tau(t))} \times \\ & \times \exp\left(\int_{\tau^\infty(x)}^t \frac{\ln \frac{B_k(\tau(s))}{(\tau(s)-\tau^\infty(s))(s-\tau^\infty(s))(s-\tau(s))(\tau(s)-\tau^2(s))\varphi_k(\tau(s))^2\eta_k(\tau(s))}}{s-\tau(s)} d_\tau s\right) d_\tau t. \end{aligned}$$

Thus, by (A.15) and (A.36) one gets

$$\begin{aligned} \xi_k(x) = & (x - \tau^\infty(x))^{-1} \exp\left(-\int_{\tau^\infty(x)}^x \frac{\ln \frac{(\tau(t)-\tau^\infty(t))(t-\tau(t))(\tau(t)-\tau^2(t))\varphi_k(\tau(t))^2\eta_k(\tau(t))}{(t-\tau^\infty(t))B_k(\tau(t))}}{t-\tau(t)} d_\tau t\right) \times \\ & \times \frac{1}{\xi_{k0}^{-1} - \int_{\tau^\infty(x)}^x \frac{(\tau(t)-\tau^2(t))}{(t-\tau^\infty(t))B_k(\tau(t))} d_\tau t} \exp\left(\int_{\tau^\infty(x)}^t \frac{\ln \frac{B_k(\tau(s))}{(\tau(s)-\tau^\infty(s))(s-\tau^\infty(s))(s-\tau(s))(\tau(s)-\tau^2(s))\varphi_k(\tau(s))^2\eta_k(\tau(s))}}{s-\tau(s)} d_\tau s\right) d_\tau t. \end{aligned} \quad (5.8)$$

From (5.8) by application of (5.1), (3.7), (3.8), (2.10), (2.11) and (3.3) we can write down explicit form of operators $\mathbf{A}_{k+1}$ and $\mathbf{A}_{k+1}^*$. We will not do it here due to length of formulas.

5.2. q -Hahn polynomials. Let us show now the well-know application of the method described in a previous section to q -deformed classical polynomials. We consider $\tau(x) = qx$ for $0 < q < 1$ and to stress it we replace the symbol ∂_τ and T by ∂_q and Q respectively. We assume that B_0 and A_0 defined by (2.12) are the second and first order polynomials respectively. We consider a function ρ_0 satisfying q -Pearson equation (2.13) and we put $f_0 \equiv 0$, $h_0 \equiv 1$. In this case the equation (3.1) is known as Hahn equation. Its solutions are given by basic hypergeometric series, see [8, 6], and we know from the theory of q -classical orthogonal polynomials that for certain choices of λ_0 , we obtain polynomials which belong to $\mathcal{H}_0$ and form orthogonal system with respect to weight function ρ_0 , see [12].

We want to keep the same type of equation for all k , i.e. we require that all $f_k \equiv 0$, $h_k \equiv 1$. We shall find the conditions under which this requirement can be satisfied along with relation (3.4).

Equation (3.8) reduces itself to

$$(5.9) \quad g_k d_k = q^{-1}.$$

The transformation rule for η_k (2.11) can be rewritten in terms of A_k as follows

$$(5.10) \quad A_{k+1}(x) = qQ A_k(x) + \partial_q B_{k+1}(x).$$

Thus A_k and B_k remain a polynomials of degree one or two respectively for all k .

We can express function ξ_k defined by (5.1) in terms of our variables as

$$(5.11) \quad \xi_k = \frac{A_k}{(1-q)x}$$

and equation (5.2) is satisfied for $c_k = -\partial_q A_k$, which is a constant for all k . Let us mention that since ξ_k is singular we can obtain this solution from (5.2) by the product (A.15) described in Appendix A only by applying singular Darboux transform (A.34) with $\delta_1 - \delta_2 = 1$.

Formula (3.5) yields that if $\{P_n^k\}$ is an OPS for ρ_k then $\{\partial_q P_n^k\}$ is an OPS for ρ_{k+1} . We can reverse this procedure. Namely, by applying $\mathbf{A}_k^*$ to both sides of (3.5) we gather that

$$(5.12) \quad \psi_k = \frac{1}{\lambda_k} \mathbf{A}_k^* \psi_{k+1}.$$

This allows us to express any polynomial by the formula

$$(5.13) \quad P_n^k = \mu_n^k \mathbf{A}_{k+1}^* \cdots \mathbf{A}_{k+n}^* 1,$$

where $\mu_n^k \neq 0$ are some normalization constants, see [12].

We can generalize this result by dropping the requirement that $f_k \equiv 0$ for $k \geq 1$ and then put $c_k = 0$. Then we can solve the equation (5.2) by the formula (5.8). We can express that solution by basic hypergeometric function. This case is presented as an example in [7].

By the application of the change of variables described in Section 4 one can obtain solutions to another family of equations with $\tau(x) \neq qx$. For example, let $\kappa(x) = e^x$. Then $\tilde{\tau}(y) = y^q$. Hahn equation transforms into

$$(5.14) \quad \tilde{\mathbf{A}}_k^* \tilde{\mathbf{A}}_k \psi_k = 0,$$

where

$$(5.15) \quad \mathbf{A}_k^* \mathbf{A}_k = \left(\frac{1-e^q}{1-q} \frac{y}{\ln y} K^{-1} A_k + \left(\frac{1-e^q}{1-q} \right)^2 \left(\partial_{\tilde{\tau}} \frac{qy^{\frac{1}{q}}}{\ln y} \right) K^{-1} B_k \right) +$$

$$+ \left(\frac{1-e^q}{1-q} \right)^2 \frac{y}{\ln y} K^{-1} B_k \partial_{\bar{\tau}} \tilde{T}^{-1} \partial_{\bar{\tau}}.$$

Its solutions are related to orthogonal polynomials by formula (4.15) and are orthogonal with respect to the scalar product with weight function

$$(5.16) \quad \tilde{\rho}_k(y) = \frac{\ln y}{y - y^q} (1-q) K^{-1} \rho_k(y).$$

5.3. The case $g_k = \text{const}$ and $\tau(x) = qx$. This example is based on the paper [5]. We assume that

$$(5.17) \quad \tau(x) = qx \quad \text{for } 0 < q < 1,$$

$$(5.18) \quad h_k \equiv 1,$$

$$(5.19) \quad g_k = q^{-2},$$

$$(5.20) \quad d_k = 1$$

and that B_0 is a constant function. From (2.10) we gather that

$$(5.21) \quad B_k = q^{-2k} B_0.$$

Let $\alpha_k := (\varphi_k)^2 \eta_k$. Equation (3.9) now takes the form

$$(5.22) \quad \alpha_k(x) = q^{-2} \alpha_k(qx) - c_k.$$

By substituting into equation (5.22) the series

$$(5.23) \quad \alpha_k(x) = x^\lambda \sum_{n \in \mathbb{Z}} a_n x^n,$$

where $\lambda \in [0, 1)$, we obtain

$$(5.24) \quad \alpha_k(x) = b_k x^{-2} - \frac{c_k}{1-q^2}$$

for any $b_k \in \mathbb{C}$. In order to preserve the transformation rules (2.11) and (3.8) we have to put $b_k = q^{4k} b_0$ and $c_k = q^{2k} c_0$. Under these assumptions the operators $\mathbf{A}_k$ and $\mathbf{A}_k^*$ take the form

$$(5.25) \quad \mathbf{A}_k = -\frac{1}{(1-q)x} Q + q^{2k} (Q^k \varphi_0),$$

$$(5.26) \quad \mathbf{A}_k^* = \frac{q^{2k} b_0 x^{-2} - \frac{c_0}{1-q^2}}{(Q^k \varphi_0)} - \frac{B_0}{q^{2k} (1-q)x} Q^{-1}.$$

Thus we have solved the factorization problem for the operators

$$(5.27) \quad \begin{aligned} & \mathbf{A}_k^* \mathbf{A}_k = \left(x^{-1} \frac{q^{2k} c_0}{\varphi_k(x) (1-q)^2 (1+q)} - x^{-3} \frac{q^{4k} b_0}{\varphi_k(x) (1-q)} \right) Q - \\ & - x^{-1} \frac{B_0 \varphi_k(q^{-1}x)}{q^2 (1-q)} Q^{-1} + x^{-2} \left(q^{4k} b_0 + \frac{B_0}{q(1-q)^2} \right) - \frac{q^{2k} c_0}{(1-q^2)}. \end{aligned}$$

The relation

$$(5.28) \quad \mathbf{A}_{k-1}^* \psi_k = 0$$

implies that $\mathbf{A}_k^* \mathbf{A}_k \psi_k = (\mathbf{A}_{k-1} \mathbf{A}_{k-1}^* - c_{k-1}) \psi_k = -c_{k-1} \psi_k$. Thus ψ_k is an eigenvector of $\mathbf{A}_k \mathbf{A}_k^*$ with the eigenvalue $-c_{k-1}$. We easily find the solution to (5.28) by the use of (A.3):

$$(5.29) \quad \psi_k = \kappa \exp \int_0^x \frac{\ln \left(\frac{(1-q)qt\alpha_k(t)}{B_k \varphi_k(qt)} \right)}{t(1-q)} d_q t,$$

where κ is a constant. Let us assume that (5.29) is convergent. We have to check if $\psi_k \in \mathcal{H}_k$, i.e. if it is square integrable with weight function ρ_k emerging from q -Pearson equation (2.8). To that end we shall calculate function $\psi_k^2 \rho_k$. Let us rewrite Pearson equation in the form

$$(5.30) \quad \frac{\alpha_k}{\varphi_k^2 B_k} = \frac{Q \rho_k}{\rho_k}.$$

From (5.28) we conclude that

$$(5.31) \quad \frac{Q \psi_k}{\psi_k} = \frac{B_k}{(1-q)q^3 x} \frac{\alpha_k}{\varphi_k}$$

and thus

$$(5.32) \quad \psi_k^2 \rho_k = \left(b_k - \frac{c_k}{1-q^2} x^2 \right) \frac{(1-q)^2 q^6}{B_k} Q(\psi_k^2 \rho_k).$$

We can rewrite this equation in the following form

$$(5.33) \quad \psi_k^2 \rho_k = \gamma \left(1 - \frac{x}{\beta} \right) \left(1 + \frac{x}{\beta} \right) Q(\psi_k^2 \rho_k),$$

where $\beta = \sqrt{\frac{b_k(1-q^2)}{c_k}}$ and $\gamma = -\frac{(1-q)^2 q^6 b_k}{B_k}$. If we request that $\gamma = 1$ by appropriate choice of b_k and B_k then we have the following solution

$$(5.34) \quad (\psi_k(x))^2 \rho_k(x) = \mu \left(-\frac{x}{\beta}; q \right)_\infty \left(\frac{x}{\beta}; q \right)_\infty,$$

where

$$(5.35) \quad (\alpha; q)_n := (1-\alpha)(1-q\alpha) \cdots (1-q^{n-1}\alpha)$$

and $\mu \in \mathbb{R} \setminus \{0\}$ is such that the scalar product is positively definite. Let us estimate the q -integral of this function.

$$(5.36) \quad \int_0^a (\psi_k(x))^2 \rho_k(x) d_q x = \mu(1-q)a \sum_{n=0}^{\infty} q^n \left(-\frac{q^n a}{\beta}; q \right)_\infty \left(\frac{q^n a}{\beta}; q \right)_\infty$$

Since $(\pm a/\beta; q)_n$ is convergent when $n \rightarrow \infty$ for $0 < q < 1$ then $(\pm a q^n/\beta; q)_\infty$ is necessarily convergent to 1. Thus it is bounded from below and above. Let us denote these lower bounds by $m_\pm$ and upper bounds by $M_\pm$. Thus

$$(5.37) \quad \left| \int_0^a (\psi_k(x))^2 \rho_k(x) d_q x \right| \leq |\mu a| \max(|M_-|, |m_-|) \max(|M_+|, |m_+|) (1-q) \sum_{n=0}^{\infty} q^n =$$

$$= |\mu a| \max(|M_-|, |m_-|) \max(|M_+|, |m_+|).$$

So this integral is finite as well as $\langle \psi_k | \psi_k \rangle_k = \int_a^b \psi_k^2 \rho_k d_q x$ and thus $\psi_k \in \mathcal{H}_k$ for any choice of limits of integration fulfilling (2.9) and constants such that $\gamma = 1$.

Thus by (3.5) we have

$$(5.38) \quad \psi_k^N = \mathbf{A}_{k+N-1} \cdots \mathbf{A}_{k+1} \mathbf{A}_k \psi_k \in \mathcal{H}_{k+N}$$

and ψ_k^N is an eigenvector of the operator $\mathbf{A}_{k+N}^* \mathbf{A}_{k+N}$ with the eigenvalue equal to

$$(5.39) \quad \lambda_k^N = - \sum_{l=k-1}^{k+N-1} c_l.$$

For example, if we put

$$(5.40) \quad \varphi_0(x) = \frac{b_0(1-q)q^2}{B_0 x e^{q^{-s} x^s}} - \frac{c_0(1-q)x}{B_0 e^{q^{-s} x^s}},$$

then

$$(5.41) \quad \psi_0(x) = \frac{\kappa(1-q^s)}{(1-q)^2} x^s.$$

Examples given above show the importance of the factorization method in the theory of the orthogonal polynomials and quantum mechanical spectral problems. In [5] the wide class of q -Schrödinger equations, which are solved by this method, is presented.

5.4. The case $g_k = \text{const}$ and τ is a fractional map. This example is similar to previous one but τ is kept arbitrary. We still consider the case when $h_k \equiv 1$ and $g_k = \text{const}$, $k \in \mathbb{N} \cup \{0\}$. From equation (3.9) one has that that case is possible for example if

$$(5.42) \quad B_k(x) = b_k(x - \tau(x))(\tau^{-1}(x) - x)$$

and

$$(5.43) \quad (\varphi_k(x))^2 \eta_k(x) = a_k$$

for some constants a_k, b_k . These conditions are compatible with transformation rules (2.11), (2.10) and (3.8) by following relations

$$(5.44) \quad a_{k+1} = \frac{g_k}{d_k^2} a_k$$

$$(5.45) \quad b_{k+1} = g_k b_k.$$

The equations (3.9) reduce to quadratic equations with constant coefficient and constants g_k are given as its roots. We will treat function φ_0 as arbitrary as well as constants a_0, b_0, c_k, d_k . In this case the equations (3.2) assume the form

$$(5.46) \quad \frac{-a_k}{\varphi_k(x)(x - \tau(x))} \psi_k(\tau(x)) + (b_k + a_k - \lambda_k) \psi_k(x) - b_k \varphi_k(\tau^{-1}(x))(\tau^{-1}(x) - x) \psi_k(\tau^{-1}(x)) = 0.$$

Let us consider the equations

$$(5.47) \quad \mathbf{A}_k \psi_k = 0.$$

It takes the form

$$(5.48) \quad \psi_k(x) = \frac{1}{(x - \tau(x)) \varphi_k(x)} \psi_k(\tau(x))$$

and

$$(5.49) \quad (\psi_k(x))^2 \rho_k(x) = \frac{b_k}{a_k} \partial_\tau \tau(x) (\psi_k(\tau(x)))^2 \rho_k(\tau(x)).$$

If we postulate the following form of the solution to (5.49)

$$(5.50) \quad (\psi_k(x))^2 \rho_k(x) = (x - \tau(x))^\lambda \alpha_k(x),$$

where $\lambda \in \mathbb{Z}$, then we get

$$(5.51) \quad \alpha_k(x) = \frac{b_k}{a_k} (\partial_\tau \tau(x))^{\lambda+1} \alpha_k(\tau(x)).$$

Necessary condition for this equation to have non-zero solution is

$$(5.52) \quad \left(\frac{b_k}{a_k} \right)^{\frac{1}{\lambda+1}} = \partial_\tau \tau(\tau^\infty(x)).$$

Let us now consider as an example the fractional map

$$(5.53) \quad \tau(x) = \frac{ax}{(a-1)x+1}$$

for $a > 0$, $a \neq 1$, which preserves the interval $(0, 1)$ and has two fixed points — 0 and 1. Let us set $\mathcal{S} = \mathcal{O}(\frac{1}{2})$. One can find formula for τ^k :

$$(5.54) \quad \tau^k(x) = \frac{a^k x}{(a^k - 1)x + 1}.$$

It is easily computed that

$$(5.55) \quad (\partial_\tau \tau)(x) = \frac{a}{(-1 + a^2)x + 1}$$

and

$$(5.56) \quad (\partial_\tau \tau)(\tau^k(x)) = a \frac{(a^k - 1)x + 1}{(a^{k+2} - 1)x + 1}.$$

Iteration of (5.51) gives us that

$$(5.57) \quad \alpha_k(x) = \begin{cases} \left(\frac{(a-1)x+1}{(-x+1)^2} \right)^{\lambda+1} \alpha_k(0) & 0 < a < 1 \\ \left(\frac{(a-1)x+1}{ax^2} \right)^{\lambda+1} \alpha_k(1) & a > 1 \end{cases}.$$

For $\lambda \neq -1$ the function $(\psi_k(x))^2 \rho_k(x)(x - \tau(x))$ has the pole at 0 or 1, while for $\lambda = -1$ it is constant. Thus $\psi_k \notin \mathcal{H}_k$ for any $k \in \mathbb{N} \cup \{0\}$. It is also possible to show that kernel of A_{k+1}^* also is empty.

We can still consider solutions to (5.47) as functions and use (3.5) to construct new solutions to (5.46). For example, let us fix

$$(5.58) \quad \varphi_0(x) = \frac{1}{x - \tau(x)}.$$

From (3.8) and assumptions of this example it follows that

$$(5.59) \quad \varphi_k(x) = \frac{1}{D_k G_k} \frac{1}{\tau^k(x) - \tau^{k+1}(x)},$$

where $G_k = g_k \cdots g_0$ and $D_k = d_k \cdots d_0$. Equations (5.47) assume the form

$$(5.60) \quad \psi_k(x) = D_k G_k \frac{\tau^k(x) - \tau^{k+1}(x)}{x - \tau(x)}.$$

Its solutions can be found by iteration and are given by

$$(5.61) \quad \psi_k(x) = \prod_{j=0}^{k-1} \frac{((a^j - 1)x + 1)((a^{j+1} - 1)x + 1)}{(x - 1)^2} \psi_k(0)$$

for $0 < a < 1$ and $(G_k D_k)^{\frac{1}{k}} a = 1$ or

$$(5.62) \quad \psi_k(x) = \prod_{j=0}^{k-1} \frac{((a^j - 1)x + 1)((a^{j+1} - 1)x + 1)}{a^{2j+1} x^2} \psi_k(1)$$

for $a > 1$ and $(G_k D_k)^{\frac{1}{k}} = a$.

From these solutions we can construct new by (3.5).

APPENDIX A. τ -RICCATI EQUATION AND τ -DARBOUX TRANSFORM

We will present here additional information about τ -difference equations and methods of solving. Most of this appendix is a direct generalization of [13].

Let us begin this section by solving the equation

$$(A.1) \quad \partial_\tau \varphi = \varphi$$

with initial condition $\varphi(\tau^\infty(x)) = 1$, under the assumption that τ has the contracting property $|\tau(x) - \tau(y)| \leq M|x - y|$, where $0 < M < 1$. Then, by the iteration one finds

$$(A.2) \quad \varphi(x) = \exp_\tau(x) := \prod_{n=0}^{\infty} \frac{1}{1 - \tau^n(x) + \tau^{n+1}(x)}.$$

We shall call $\exp_\tau$ the τ -**exponential** function.

Let us formulate one more identity which was used throughout the paper.

$$(A.3) \quad \prod_{n=0}^{\infty} \left(T^n F(x) \right) = \exp \left(\int_{\tau^\infty(x)}^x \frac{\ln F(t)}{t - \tau(t)} d_\tau t \right).$$

It follows from the observation that

$$(A.4) \quad \ln \prod_{n=0}^{\infty} \left(T^n F(x) \right) = \sum_{n=0}^{\infty} T^n \ln F(x)$$

and from the definition of τ -integral (1.9). Let us note that $\exp$ and $\ln$ appearing in (A.3) are standard exponent and logarithm.

From (A.3) it follows that the solution to

$$(A.5) \quad \partial_\tau \psi = f\psi$$

is given by

$$(A.6) \quad \psi(x) = \exp \left(\int_{\tau^\infty(x)}^x \frac{-\ln(1 - (t - \tau(t))f(t))}{t - \tau(t)} d_\tau t \right) \psi(\tau^\infty(x)).$$

Now, we are going to investigate the functional equation of the form

$$(A.7) \quad \begin{pmatrix} \partial_\tau \psi(x) \\ \partial_\tau \varphi(x) \end{pmatrix} = \tilde{\mathbf{A}}(x) \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix} = \begin{pmatrix} \tilde{a}(x) & \tilde{b}(x) \\ \tilde{c}(x) & \tilde{d}(x) \end{pmatrix} \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix}.$$

By the substitution

$$(A.8) \quad \mathbf{A}(x) = \mathbb{I} - (x - \tau(x))\tilde{\mathbf{A}}(x)$$

it can be equivalently written as

$$(A.9) \quad \begin{pmatrix} T\psi(x) \\ T\varphi(x) \end{pmatrix} = \mathbf{\Lambda}(x) \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix} = \begin{pmatrix} a(x) & b(x) \\ c(x) & d(x) \end{pmatrix} \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix}.$$

If we introduce the function

$$(A.10) \quad u(x) := \frac{\varphi(x)}{\psi(x)},$$

the equation (A.9) assumes the form

$$(A.11) \quad u(\tau(x)) = \frac{d(x)u(x) + c(x)}{b(x)u(x) + a(x)},$$

which is equivalent to

$$(A.12) \quad \partial_\tau u(x) = \tilde{c}(x) + \tilde{d}(x)u(x) - \tilde{a}(x)u(\tau(x)) - \tilde{b}(x)u(x)u(\tau(x)).$$

Equation (A.12) is generalization of q -Riccati equation and thus we will call both (A.11) and (A.12) the **τ -Riccati equation**.

Equation (3.1) considered in Section 3 can be obtained from (A.9) by putting

$$(A.13) \quad \mathbf{\Lambda}(x) = \begin{pmatrix} \frac{\lambda - \beta(x)}{\alpha(x)} & -\frac{\gamma(x)}{\alpha(x)} \\ 1 & 0 \end{pmatrix}.$$

The formal solution to (A.9) is given in terms of the infinite matrix product

$$(A.14) \quad \mathbf{\Lambda}_\infty(x) := \lim_{n \rightarrow \infty} \mathbf{\Lambda}(\tau^n(x)) \cdots \mathbf{\Lambda}(\tau^2(x)) \mathbf{\Lambda}(\tau(x)) \mathbf{\Lambda}(x)$$

by the formula

$$(A.15) \quad \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix} = \mathbf{\Lambda}(x)_\infty^{-1} \begin{pmatrix} \psi(\tau^\infty(x)) \\ \varphi(\tau^\infty(x)) \end{pmatrix}.$$

Sometimes we are able to compute resolvent function (A.14) explicitly.

Proposition A.1.

If a , b and d are continuous at $\tau^\infty(x)$ and $c(x) \equiv 0$ then

$$(A.16) \quad \mathbf{\Lambda}_\infty(x) = \begin{pmatrix} \exp \int_{\tau^\infty(x)}^x \frac{\ln a(t)}{t - \tau(t)} d_\tau t & F(x) \\ 0 & \exp \int_{\tau^\infty(x)}^x \frac{\ln d(t)}{t - \tau(t)} d_\tau t \end{pmatrix},$$

where $F(x)$ is given by

$$(A.17) \quad F(x) = \exp \left(\int_{\tau^\infty(x)}^x \frac{\ln d(t)}{t - \tau(t)} d_\tau t \right) \int_{\tau^\infty(x)}^x \frac{b(t)}{(t - \tau(t))a(t)} \exp \left(\int_{\tau^\infty(x)}^t \frac{\ln \frac{a(s)}{d(s)}}{s - \tau(s)} d_\tau s \right) d_\tau t$$

Proof: From the fact that $\mathbf{\Lambda}$ is upper-triangular we conclude that $\mathbf{\Lambda}_\infty$ is given by

$$(A.18) \quad \mathbf{\Lambda}_\infty(x) = \begin{pmatrix} a_\infty(x) & F(x) \\ 0 & d_\infty(x) \end{pmatrix},$$

where $a_\infty(x) = \prod_{n=0}^\infty T^n a(x)$ and $d_\infty(x) = \prod_{n=0}^\infty T^n d(x)$. Since

$$(A.19) \quad \mathbf{\Lambda}_\infty(x) = \mathbf{\Lambda}_\infty(\tau(x)) \mathbf{\Lambda}(x)$$

we see that $F(x)$ satisfies the equation

$$(A.20) \quad F(x) = d(x)F(\tau(x)) + b(x)d_\infty(\tau(x)).$$

Solving (A.20) we get

$$(A.21) \quad F(x) = \left(\prod_{l=0}^{\infty} d(\tau^l(x)) \right) \sum_{k=0}^{\infty} \frac{b(\tau^k(x))}{a(\tau^k(x))} \prod_{j=k}^{\infty} \frac{a(\tau^j(x))}{d(\tau^j(x))}.$$

From (A.3), (A.6) and (1.8) we obtain now (A.17). $\square$

We should note that if one considers equations (A.11) or (A.12) then the matrix-valued function $\mathbf{\Lambda}$ is defined up to a function factor and this observation can be used to regularize the product (A.14).

We are going to consider the following gauge-like transformation

$$(A.22) \quad \mathbf{\Lambda}(x) \longrightarrow \mathbf{\Lambda}'(x) = D(\tau(x))^{-1} \mathbf{\Lambda}(x) D(x),$$

for some $D : X \rightarrow GL(2, \mathbb{C})$, which transforms the functional equation (A.9) into the one of the same type with matrix $\mathbf{\Lambda}'$. Its solutions are given by

$$(A.23) \quad \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix} \longrightarrow D(x)^{-1} \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix}.$$

The resolvent of new equation is given by

$$(A.24) \quad \mathbf{\Lambda}_{\infty}(x) \longrightarrow D(0)^{-1} \mathbf{\Lambda}_{\infty}(x) D(x).$$

We call this transform the τ -**Darboux transform**. It is a generalization of q -Darboux transform introduced in [13]. Sometimes we are able to reduce $\mathbf{\Lambda}$ to such $\mathbf{\Lambda}'$ that we can compute (A.14) explicitly. This is the case, for example, if $\mathbf{\Lambda}'$ is upper-triangular. Applying this method, we can obtain the result for τ -Riccati case, which generalize this of q -Riccati equation of [13].

This is a classical result for the Riccati equation generalized to q -case in [13] and here it is presented in τ -version.

Proposition A.2.

Let u_0 be a particular solution of (A.11). The general solution of (A.9) is then given by the formula

$$(A.25) \quad \begin{aligned} \psi(x) = & \exp \left(- \int_{\tau^{\infty}(x)}^x \frac{\ln a(t) + b(t)u_0(t)}{t - \tau(t)} d_{\tau} t \right) \left(-B \int_{\tau^{\infty}(x)}^x \frac{b(t)}{a(t) + b(t)u_0(t)} \times \right. \\ & \times \exp \left(\int_{\tau^{\infty}(x)}^t \frac{\ln \frac{a(s) + b(s)u_0(s)}{-b(s)u_0(\tau(s)) + d(s)}}{s - \tau(s)} d_{\tau} s \right) d_{\tau} t + A \Big) \end{aligned}$$

$$(A.26) \quad \varphi(x) = u_0(x)\psi(x) + B \exp \left(- \int_{\tau^{\infty}(x)}^x \frac{\ln -b(t)u_0(\tau(t)) + d(t)}{t - \tau(t)} d_{\tau} t \right),$$

where the constants A and B are related to the initial conditions in the following way

$$(A.27) \quad A = \psi(\tau^{\infty}(x))$$

and

$$(A.28) \quad B = -\psi(\tau^{\infty}(x))u_0(\tau^{\infty}(x)) + \varphi(\tau^{\infty}(x)).$$

Solution of (A.11) is then given by
(A.29)

$$u^t(x) = u_0(x) + \frac{t \exp \left(\int_{\tau^\infty(x)}^x \frac{\ln \frac{a(z)+b(z)u_0(z)}{-b(z)u_0(\tau(z))+d(z)}}{z-\tau(z)} d_\tau z \right)}{1 - t \int_{\tau^\infty(x)}^x \frac{b(z)}{(z-\tau(z))(a(z)+b(z)u_0(z))} \exp \left(\int_{\tau^\infty(x)}^z \frac{\ln \frac{a(s)+b(s)u_0(s)}{-b(s)u_0(\tau(s))+d(s)}}{s-\tau(s)} d_\tau s \right) d_\tau z}$$

for any $t \in \mathbb{R}$.

Proof: In order to obtain the general solution of (A.7), we apply the τ -Darboux transform (A.22) with $D(x)$ of the form

$$(A.30) \quad D(x) = \begin{pmatrix} 1 & 0 \\ u_0(x) & 1 \end{pmatrix}.$$

Since u_0 is a particular solution of (A.11), the transformed matrix $\Lambda'(x)$ is upper-triangular. Solution given by (A.2) and (A.26) is obtained by substituting into (A.15) the formulae (A.16) and (A.17) from Proposition A.1 and transforming formula (A.24) for resolvent. Formula (A.29) follows from (A.10), (A.2) and (A.26) by putting $t := \frac{B}{A}$. $\square$

We define the transform $\mathcal{B}_t^+$ by

$$(A.31) \quad \mathcal{B}_t^+ u_0 := u^t.$$

It is easy to check that $\mathcal{B}_t^+$, $t \in \mathbb{R}$, form a one-parameter group

$$(A.32) \quad B_s^+ \circ B_t^+ = B_{s+t}^+.$$

Moreover the solutions $u^{t_1}(x)$, $u^{t_2}(x)$, $u^{t_3}(x)$ and $u^{t_4}(x)$ do satisfy the unharmonical superposition principle:

$$(A.33) \quad \frac{(u^{t_4}(x) - u^{t_3}(x))(u^{t_1}(x) - u^{t_2}(x))}{(u^{t_3}(x) - u^{t_1}(x))(u^{t_2}(x) - u^{t_4}(x))} = \frac{(t_4 - t_3)(t_1 - t_2)}{(t_3 - t_1)(t_2 - t_4)}.$$

The proof of (A.32) and (A.33) is straightforward.

Let us mention another application of τ -Darboux transform. When $\Lambda(\tau^\infty(x)) = \mathbb{I}$ we call (A.9) the regular equation. Solutions of regular equation are finite and nonzero at $\tau^\infty(x)$. Let us introduce singularity (or zero) at this point by the τ -Darboux transform (A.22) - (A.23) with

$$(A.34) \quad D(x) = \begin{pmatrix} (x - \tau^\infty(x))^{\delta_1} & 0 \\ 0 & (x - \tau^\infty(x))^{\delta_2} \end{pmatrix}.$$

Thus one has

$$(A.35) \quad \Lambda'(x) = \begin{pmatrix} a(x) \left(\frac{x - \tau^\infty(x)}{\tau(x) - \tau^\infty(x)} \right)^{\delta_1} & b(x) \frac{(x - \tau^\infty(x))^{\delta_2}}{(\tau(x) - \tau^\infty(x))^{\delta_1}} \\ c(x) \frac{(x - \tau^\infty(x))^{\delta_1}}{(\tau(x) - \tau^\infty(x))^{\delta_2}} & d(x) \left(\frac{x - \tau^\infty(x)}{\tau(x) - \tau^\infty(x)} \right)^{\delta_2} \end{pmatrix}$$

and the solution to (A.11) with $\Lambda'(x)$ is

$$(A.36) \quad u'(x) = (x - \tau^\infty(x))^{\delta_1 - \delta_2} u(x),$$

where $u(x)$ is a solution to (A.11) with $\Lambda(x)$.

Hence one can have $u'(\tau^\infty(x)) = \infty$ for $\delta_1 - \delta_2 < 0$ while $u(\tau^\infty(x)) < \infty$. Thus we can apply the transformation (A.34) to regularize the singular equation of the type (A.35) to the regular one. It was applied implicitly for example in (5.50).

Infinite products in previous formulae were always assumed to be convergent. We will give now a condition for this to be the case.

Proposition A.3.

Let the following conditions be satisfied:

- i) $\sum_{k=0}^{\infty} |\tau^k(x) - \tau^{k+1}(x)| < \infty$, e.g. it has a place if τ is a contracting map;
- ii) $\tilde{\mathbf{A}}$ is continuous at $\tau^\infty(x)$.

Then product (A.14) is convergent.

Proof: We will show that under these assumptions $P_n = \mathbf{A}(\tau^n(x)) \cdots \mathbf{A}(x)$ is a Cauchy sequence and thus convergent. We can estimate the norm of the difference between two terms of this series by

$$(A.37) \quad \|P_n - P_{n+k}\| \leq \left\| \mathbf{A}(\tau^{n+k}(x)) \cdots \mathbf{A}(\tau^{n+1}(x)) - 1 \right\| \|P_n\|.$$

Let us show that it can be made arbitrarily small. To that end we show first that $\|P_n\|$ is bounded.

$$(A.38) \quad \|P_n\| \leq \prod_{k=0}^n (1 + |\tau^k(x) - \tau^{k+1}(x)| \left\| \tilde{\mathbf{A}}(\tau^k(x)) \right\|).$$

This product is convergent if and only if

$$(A.39) \quad \sum_{n=0}^{\infty} |\tau^n(x) - \tau^{n+1}(x)| \left\| \tilde{\mathbf{A}}(\tau^n(x)) \right\| < \infty.$$

Assumptions of the proposition guarantees that this is the case. Hence

$$(A.40) \quad \|P_n\| \leq \prod_{k=0}^{\infty} (1 + |\tau^k(x) - \tau^{k+1}(x)| \left\| \tilde{\mathbf{A}}(\tau^k(x)) \right\|) < \infty.$$

Now we show that for n big enough $\left\| \mathbf{A}(\tau^{n+k}(x)) \cdots \mathbf{A}(\tau^{n+1}(x)) - 1 \right\|$ is arbitrarily small. We have the inequality

$$(A.41) \quad \left\| \mathbf{A}(\tau^{n+k}(x)) \cdots \mathbf{A}(\tau^{n+1}(x)) - 1 \right\| + 1 \leq \\ \leq \left(1 + |\tau^{n+k}(x) - \tau^{n+k+1}(x)| \left\| \tilde{\mathbf{A}}(\tau^{n+k}(x)) \right\| \right) \cdots \left(1 + |\tau^{n+1}(x) - \tau^{n+2}(x)| \left\| \tilde{\mathbf{A}}(\tau^{n+1}(x)) \right\| \right).$$

Since (A.39) hold then (A.41) tends to 1 as $n \rightarrow \infty$. Thus we have completed the proof that P_n is Cauchy sequence. $\square$

ACKNOWLEDGEMENT

Authors would like to thank dr. M. Klimek for the information about works [3, 11] and dr. S. Stepin, dr. M. Horowski for careful reading of the manuscript and interest in the paper. We would also like to thank Alina Dobrogowska for help and fruitful discussions. This work is supported in part by KBN grant 2 PO3 A 012 19.

REFERENCES

- [1] N. I. Akhiezer and I. M. Glazman. *Theory of Linear Operators in Hilbert Space*. Dover Publications, 1993.
- [2] A. Antonevich. *Linear Functional Equations - Operator Approach*. Birkhauser Verlag AG, 1995.
- [3] A. Arai. Exactly solvable supersymmetric quantum mechanics. *J. Math. Anal. Appl.*, 158:63–79, 1991.
- [4] G. Darboux. Sur une proposition relative aux equations lineaires. *C. R. Acad. Sci. Paris*, 94:1456, 1882.
- [5] A. Dobrogowska and A. Odziejewicz. Second order q -difference equations solvable by factorization method. *math-ph/0312057*, 2003.
- [6] G. Gasper and M. Rahman. *Basic Hypergeometric Series*. Cambridge University Press, 1990.
- [7] T. Goliński and A. Odziejewicz. General difference calculus and its application to functional equations of the second order. *Czech. J. Phys.*, 52:1219–1218, 2002.
- [8] W. Hahn. Über Orthogonalpolynome die q -Differenzengleichungen genügen. *Math. Nachr.*, 2:4–34, 1949.
- [9] M. Horowski, A. Odziejewicz, and A. Tereszkiewicz. Some integrable systems in nonlinear quantum optics. *Journal of Mathematical Physics*, 44(2):480–506, 2002.
- [10] L. Infeld and T. E. Hull. The factorization method. *Rev. Mod. Phys.*, 23:21, 1951.
- [11] M. Klimek. Extension of q -deformed analysis and q -deformed models of classical mechanics. *J. Phys. A: Math. Gen.*, 26:955–967, 1993.
- [12] A. Odziejewicz, M. Horowski, and A. Tereszkiewicz. Integrable multi-boson systems and orthogonal polynomials. *J. Phys. A: Math. Gen.*, 34:4353–4376, 2001.
- [13] A. Odziejewicz and A. Ryżko. The Darboux-like transform and some integrable cases of the q -Riccati equation. *J. Phys. A: Math. Gen.*, 35:747–757, 2002.